# Cellwet®

# Eve Drops

# (Polyethylene Glycol 400 & Propylene Glycol Ophthalmic Solution (0.4% & 0.3% w/v))

### COMPOSITION:

Polyethylene Glycol 400 IP	0.4% w/v
Propylene Glycol IP	0.3% w/v
Sodium Perborate BP	005% w/\
(as preservative)	
Water for Injections IP	q.s.

### PHARMACOLOGICAL CLASSIFICATION:

Lubricant

# PHARMACOLOGY:

Biphasic mechanism of action in which the product first binds to damaged hydrophobic areas of epithelial cells to add volume to the tear film and then restructures the tear film by forming a protective gel matrix that provides long lasting protection. The high molecular weight of Hydroxypropyl guar molecules, along with the natural mucin layer, prolong retention of the eye drops which provides sustained lubrication to the eye and protects the ocular surface from further damage while the surface epithelial cells undergo repair and renewal.

### INDICATIONS:

CELLWET is indicated for the temporary relief of burning and irritation due to dryness of the eye.

### CONTRA-INDICATIONS:

CELLWET is contraindicated in patients hypersensitivity to any of the ingredients.

# WARNINGS AND PRECAUTIONS:

- If irritation pain, redness or changes in vision occur or if the patients condition worsens, treatment discontinuation should be considered.
- · Remove contact lenses before using this eye drops.
- . To avoid contamination do not touch tip of container to any surface.
- · Replace the cap after using.
- . Do not use if solution changes color or becomes cloudy.

# ADVERSE EFFECTS:

CELLWET Eye Drops rarely causes serious side effects or any allergic reaction. Minor burning, stinging or irritation may temporarily occur.

### DOSAGE AND DIRECTIONS FOR USE:

Instill 1 to 2 drops in the affected eye(s) as needed.

## STORAGE:

Keep in a cool place. Use the solution within one month after opening the container.

Do not touch the tip of the container to any surface.

Protect from light

# KEEP OUT OF REACH OF CHILDREN

# NOT FOR INJECTION

### FOR EXTERNAL USE ONLY

### PRESENTATION:

CELLWET is a sterile solution supplied in plastic opaque dropper bottle with a cap, containing 10ml of the solution.

## Directions for use :



Turn the tamper proof cap anti-clockwise to break the seal. Remove the cap, dispense drops with gentle pressure.

Replace the cap immediately after every use.