

# FLI DROPS<sup>TM</sup>

## Eye Drops

(Fluorometholone Acetate  
Ophthalmic Suspension 0.1% w/v)

### COMPOSITION:

Fluorometholone Acetate USP.....	0.1% w/v
Benzalkonium chloride Solution IP .....	0.02% v/v
(As Preservative)	
Water for Injections IP .....	q.s.

### Description:

FLI DROPS contains 0.1% w/v fluorometholone acetate in sterile ophthalmic suspension.

### Uses:

This medication is used to treat inflammation and swelling in the eye caused by allergies, infections, injuries or other conditions.

### How to use:

For best effects, this medication must be used as directed. Do not stop using this medication without consulting your doctor. Remove contact lenses before using this medication. May reinsert lenses after waiting 15 minutes. To apply eye drops, wash hands first. To avoid contamination, be careful not to touch the dropper or let it touch your eye. Shake the eye drops well before using. Tilt your head back, gaze upward and pull down the lower eyelid to make a pouch. Place dropper directly over eye and administer the prescribed number of drops. Look downward and gently close your eye for 1 to 2 minutes. Place one finger at the corner of the eye near the nose and apply gentle pressure. This will prevent the medication from draining away from the eye. Try not to blink and do not rub the eye. Do not rinse the dropper. If you are using another kind of eye medication, wait at least five minutes before applying other medications.

### Side effects:

This medication may temporarily sting your eyes for a minute or two when applied. If stinging continues or becomes bothersome, inform your doctor. Vision may be temporarily blurred or unstable after applying eye drops. Use caution if driving or performing duties requiring clear vision. Notify your doctor immediately if you develop: skin rash, itching/redness/pain around the eyes, vision problems. If you notice other effects not listed above, contact your doctor or pharmacist.

### Precautions:

Tell your doctor if you have any of the following conditions: eye problems, any allergies. This medication should be used only if clearly needed during pregnancy. Discuss the risks and benefits with your doctor. It is not known if this medication is found in breast milk. Consult your doctor before breast-feeding.

### Drug interactions:

Tell your doctor of any over-the-counter or prescription medication you may take. Do not start or stop any medicine without doctor or pharmacist approval.

### Overdose:

If overdose is suspected, contact your Physician.

**Notes:**

If the condition for which this was prescribed does not improve or becomes worse after a few days, consult your doctor.

**Missed dose:**

If you miss a dose, use it as soon as remembered; do not use it if it is near the time for the next dose, instead, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

**Storage and handling instructions:**

Keep in cool place

Use the suspension within one month after opening the container.

Do not touch the tip of the container to any surface. Replace cap after use.

Protect from light.

**KEEPOUT OF REACH OF CHILDREN**

**NOT FOR INJECTION**

**FOR EXTERNAL USE ONLY**

**SHAKE WELL BEFORE USE**

**Presentation:**

FLI Drops is a sterile ophthalmic suspension supplied in opaque plastic dropper bottle with a cap, Containing 5 ml of the suspension.

**Directions for use:**

Turn the tamper proof cap anti-clockwise to break the seal.



Remove the cap, dispense drops with gentle pressure.



Replace the cap immediately after every use.

Manufactured in INDIA by :

**Senses Pharmaceuticals Pvt. Ltd.,**

No.77, 3rd Road, Bommasandra Industrial Area,  
Bommasandra 4th Phase, Bengaluru - 560 099.

TM: Trademark for Registration

Month of publication of pack insert: September 2021

FLDPI-03