Moxifix®DM

Eve Drops

Moxifloxacin with Dexamethasone Eye Drops

COMPOSITION:

Moxifloxacin Hydrochloride IP equivalent to Moxifloxacin	0.5%	w/v
Dexamethasone Sodium Phosphate IP equivalent to		
Dexamethasone Phosphate	0.1%	w/v
Water for Injections IP		q.s.

Description

Moxifix DM is a combination of Moxifloxacin 0.5% w/v and Dexamethasone 0.1% w/v. Moxifloxacin is a fourth generation fluoroquinolone controls infection by inhibiting the dna gyrase & topoisomerase iv.

Dexamethasone a potent corticosteroid effectively controls the inflammation by inhibiting the release of inflammatory mediators.

Indications

Moxifix DM is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists. The combination can also be used for post-operative inflammation and any other ocular inflammation associated with infection.

Dosage

One or two drops instilled into the conjunctival sac(s), every 4 to 6 hours. During the initial 24 to 48 hours, the dosage may be increased to 1 or 2 drops every two hours. Frequency must be decreased gradually or warranted by improvement in clinical signs. Care should be taken not to discontinue the therapy prematurely.

Contraindications

Moxifix DMis contraindicated in epithelial herpes simplex keratitis (Dendritic keratitis), vaccinia, varicella, and in many other viral diseases of the conjunctiva and cornea, Mycobacterial infection of the eye and fungal diseases of ocular structures and in individuals hypersensitive to any of the components of the medication.

Warnings & precautions

Prolonged use of steroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and posterior subcapsular cataract formation.

Adverse effects

The most frequently reported drug-related undesirable effects seen with moxifloxacin are conjunctival irritation, increased lacrimation, keratitis and papillary conjunctivitis.

Storage

Store Below 30°C, Protect from light.

DO NOT FREEZE

KEEP OUT OF REACH OF CHILDREN

NOT FOR INJECTION

FOR EXTERNAL USE ONLY

Solution to be used within one month after opening the container.

Presentation

Moxifix DM eye drops a sterile greenish yellow coloured solution is available in opaque plastic dropper bottle with a cap containing 10mL of the solution

Directions for use:



Turn the tamper proof cap anti clockwise to break the seal and remove the cap.

Remove and discard the tamper proof ring from the container.

Tilt your head back. Gently pull your lower eyelid downwards to form a pocket between your eyelid and your eye, look up.



Turn the container upside down, place the dropper tip close to your eye but be careful not to touch your eye with it and gently press the container with thumb and index finger to dispense the drop.

Replace the cap by turning clockwise direction until it is firmly

touching the bottle. Do not overtighten the cap.

Manufactured in INDIA by :