

# Naphace<sup>®</sup>

## Eye Drops

### Naphazoline Hydrochloride + Chlorpheniramine Maleate + Methyl Cellulose eye drops

For Ophthalmic Use Only

#### COMPOSITION:

Naphazoline Hydrochloride USP .....	0.1% w/v
Chlorpheniramine Maleate IP .....	0.01% w/v
Methyl Cellulose IP .....	0.1% w/v
Preservative:	
Phenyl Mercuric Nitrate IP .....	0.001%w/v
Sterile Isotonic Aqueous base .....	q.s.

#### Pharmacological properties:

Naphace eye drops as an anti-allergic, astringent & decongestant.

Naphazoline acts as a vasoconstrictor to narrow swollen blood vessels to reduce the redness.

Methyl cellulose acts as lubricant to give relief from dry irritated eyes.

Chlorpheniramine maleate acts as anti-allergic gives relief from itching, watering of eyes due to allergy.

#### Indications:

Naphace eye drops is an ophthalmic medicine used to treat allergic conjunctivitis (eye allergy).

It is indicated for temporary relief of redness and discomfort due to minor eye irritations and relief of burning, watery eyes and irritation due to dryness of the eye or exposure to wind, sun, or dust.

#### Dosage and Administration:

Instill one or two drops in the affected eye(s) 3 times a day or use eye drops as instructed by your doctor.

Wash your hands thoroughly before and after applying. Use the exact number of drops recommended.

After instilling the drops, close your eyes for 2-3 minutes; do not blink or squeeze.

Avoid touching your eyes with the dropper, it must be kept clean.

Wipe any excess liquid from your face with a napkin or tissue.

If you are prescribed any other eye drops, you need to keep an interval of at least 5 minutes between both the drops.

#### Warning and Precautions:

NOT FOR INJECTION. FOR OPHTHALMIC USE ONLY .

To avoid contamination do not touch tip of container to any surface. Shake well before use and Replace cap after using. If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the conditions worsens or persists for more than 72 hours, discontinue use and consult a doctor.

**Adverse Effects:**

Transient burning pain, and irritation stinging upon instillation and other minor symptoms of ocular. Blurred vision

**Drug Interactions:**

**Drug-Drug Interaction:** Naphazolin eye drops may interact with medicines used to treat hypertension (atenolol, propranolol)

**Drug-Food Interaction:** cause increased drowsiness, dizziness, and lack of concentration.

**Overdose:**

Naphazolin eye drops is meant for external application, cases of overdose are unlikely. However, upon accidental ingestion, contact your doctor as soon as possible or visit the nearest hospital.

**Storage and Handling Instructions:**

Keep in a cool place. Use the solution within one month after opening the container. Do not touch the tip of the container to any surface. Replace cap after using. Protect from light.

**KEEP OUT OF REACH OF CHILDREN**

**NOT FOR INJECTION**

**FOR EXTERNAL USE ONLY**

**Presentation:**

Naphase is a sterile solution supplied in white opaque plastic dropper bottle with a cap, Containing 5 ml of the solution.

**Directions for use :**

Turn the tamper proof cap anti-clockwise to break the seal.

Remove the cap, dispense drops with gentle pressure.

Replace the cap immediately after every use.

Manufactured by:

**Divine Laboratories Pvt. Ltd.**

Block No. 471, Dabhasa, Tal-Padra,

Dist- Vadodara - 391 440, India.

Marketed by:

**Senses Pharmaceuticals Pvt. Ltd.,**

#01, Survey No. 29/1&2, Yelachanahalli,  
Chunchagatta Main Road, Bangalore - 560 062, India.

Email: [info@sensespharma.com](mailto:info@sensespharma.com)

® Registered Trademark