SENZMOX®-DM

Eve Drops

Moxifloxacin with Dexamethasone Eye Drops

COMPOSITION:

| Moxifloxacin Hydrochloride IP equivalent to Moxifloxacin 0 | .5% | w/v |
|--|-------|------|
| Dexamethasone Sodium Phosphate IP equivalent to | | |
| Dexamethasone Phosphate 0. | .1% ۱ | w/v |
| Water for Injections IP | c | Į.S. |

Description

Senzmox-DM is a combination of Moxifloxacin 0.5% w/v and Dexamethasone 0.1% w/v. Moxifloxacin is a fourth generation fluoroquinolone controls infection by inhibiting the dna gyrase & topoisomerase iv.

Dexamethasone a potent corticosteroid effectively controls the inflammation by inhibiting the release of inflammatory mediators.

Indications

SENZMOX-DM is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists. The combination can also be used for post-operative inflammation and any other ocular inflammation associated with infection.

Dosage

One or two drops instilled into the conjunctival sac(s), every 4 to 6 hours. During the initial 24 to 48 hours, the dosage may be increased to 1 or 2 drops every two hours. Frequency must be decreased gradually or warranted by improvement in clinical signs. Care should be taken not to discontinue the therapy prematurely.

Contraindications

SENZMOX-DM is contraindicated in epithelial herpes simplex keratitis (Dendritic keratitis), vaccinia, varicella, and in many other viral diseases of the conjunctiva and cornea, Mycobacterial infection of the eye and fungal diseases of ocular structures and in individuals hypersensitive to any of the components of the medication.

Warnings & precautions

Prolonged use of steroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and posterior subcapsular cataract formation.

Adverse effects

The most frequently reported drug-related undesirable effects seen with moxifloxacin are conjunctival irritation, increased lacrimation, keratitis and papillary conjunctivitis.

Storage

Store in a cool dry place. Protect from light.

DO NOT FREEZE

KEEP OUT OF REACH OF CHILDREN

NOT FOR INJECTION

FOR EXTERNAL USE ONLY

Solution to be used with in one month after opening the container.

Presentation

SENZMOX-DM eye drops is available in sterile opaque plastic dropper bottle with a cap containing 5ml of the solution

Directions for use :



Turn the tamper proof cap anti-clockwise to break the seal. Remove the cap, dispense drops with gentle pressure.



Replace the cap immediately after every use.