

<sup>Rx</sup>  
**Travasenz<sup>®</sup> T**  
Eye Drops 5 mL  
(Travoprost & Timolol Eye Drops)

**Composition:**

Travoprost IP .....	0.004% w/v
Timolol Maleate IP equivalent to Timolol .....	0.5% w/v
Benzalkonium Chloride Solution IP .....	0.02% v/v
(as preservative)	
Water for Injections IP.....	q.s.

**Description:**

Travasenz T eye drops is a combination of two active substances (Travoprost and Timolol). Travoprost is a prostaglandin analogue which works by increasing the outflow of aqueous fluid from the eye, which lowers its pressure. Timolol is a beta-blocker which works by reducing the production of fluid within the eye. The two substances work together to reduce pressure within the eye.

**Indication:**

Travasenz T eye drops is indicated for the decrease of intraocular pressure (IOP) in adult patients with open angle glaucoma or ocular hypertension who are insufficiently responsive to topical beta blockers or prostaglandin analogues.

**Dosage regimen and administration:**

**General target population:**

**Adults:**

The recommended dosage of Travasenz T is one drop in the conjunctival sac of the affected eye(s) once daily in the morning or evening or as directed by the Ophthalmologist. It should be administered at the same time each day.

The dosage of Travasenz T eye drops should not exceed one drop in the affected eye(s) once daily since it has been shown that more frequent administration of prostaglandin analogues may decrease the IOP lowering effect.

If a dose is missed, treatment should be continued with the next dose as normal. If more than one topical ophthalmic drug is being used, the drugs should be administered at least 5 minutes apart.

**Children and adolescents:**

Travasenz T is not to be used by children and adolescents under 18 years of age.

**Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Do not use Travasenz T, if you are pregnant, unless your doctor considers it necessary. If you could get pregnant, you must use adequate contraception, whilst you use the medicine.

Do not use Travasenz T if you are breast-feeding. This medicine may get into your milk.

**Warnings and precautions:**

Talk to your doctor before using Travasenz T, if you have now or have had in the past

- Coronary heart disease (symptoms can include chest pain or tightness, breathlessness or choking), heart failure, low blood pressure.
- Disturbances of heart rate such as slow heartbeat.
- Breathing problems, asthma or chronic obstructive pulmonary disease.
- Poor blood circulation disease (such as Raynaud's disease or Raynaud's syndrome).
- Diabetes (as timolol may mask signs and symptoms of low blood sugar).
- Overactivity of the thyroid gland (as timolol may mask signs and symptoms of thyroid disease).
- Myasthenia gravis (chronic neuromuscular weakness).
- Cataract surgery.
- Eye inflammation.

If you need to have any type of surgery, tell your doctor that you are using Travasenz T, as timolol may change the effects of some medicines used during anaesthesia.

If you get any severe allergic reaction (skin rash, redness and itching of the eye) while using Travasenz T, whatever the cause, adrenaline treatment may not be as effective. It is therefore important to tell the doctor that you are using Travasenz T, when you are to receive any other treatment.

Travasenz T may change the colour of your iris (the coloured part of your eye). This change may be permanent.

Travasenz T may increase the length, thickness, colour and/or number of your eyelashes and may cause unusual hair growth on your eyelids.

Travoprost may be absorbed through the skin and therefore should not be used by women who are pregnant or are attempting to become pregnant. If any of the medicine comes into contact with the skin then it should be washed off straight away.

#### **Other medicines and Travasenz T**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Travasenz T can affect or be affected by other medicines you are using, including other eye drops for the treatment of glaucoma. Tell your doctor if you are using or intend to use

- Medicines to lower blood pressure,
- Heart medicines including quinidine (used to treat heart conditions and some types of malaria),
- Medicines to treat diabetes or the antidepressants fluoxetine or paroxetine.

#### **Driving and using machines**

You may find that your vision is blurred for a time just after you use Travasenz T. Do not drive or use machines until this has worn off.

Benzalkonium chloride may be absorbed by soft contact lenses and may change the colour of the contact lenses. You should remove contact lenses before using this medicine and put them back in 15 minutes afterwards.

Benzalkonium chloride may also cause eye irritation, especially if you have dry eyes or disorders of the cornea (the clear layer at the front of the eye). If you feel abnormal eye sensation, stinging or pain in the eye after using this medicine, talk to your doctor.

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### **Storage:**

Keep in cool place. Use the solution within one month after opening the container.

#### **PROTECT FROM LIGHT**

#### **KEEP OUT OF REACH OF CHILDREN**

#### **NOT FOR INJECTION**

#### **FOR EXTERNAL USE ONLY**

#### **Presentation:**

**Travasenz T** is a sterile ophthalmic solution supplied in a sterile plastic opaque dropper bottle with a cap containing 5 mL of the solution

#### **Directions for use :**



Turn the tamper proof cap anti-clockwise to break the seal.



Remove the cap, dispense drops with gentle pressure.



Replace the cap immediately after every use.

Manufactured in INDIA by :

**Senses Pharmaceuticals Pvt. Ltd.,**

No.77, 3rd Road, Bommasandra Industrial Area,  
Bommasandra 4th Phase, Bengaluru - 560 099.

© Registered Trademark

TVTRI-01

Month of publication of pack insert: February 2024